Author Index Vol. 782

Abidi, S.L. and Mounts, T.L.
Reversed-phase high-performance liquid chromatographic separations of tocopherols 782(1997)25
Affonso, R., see Dalmora, S. 782(1997)199
Anderson, J.M., see Saari-Nordhaus, R. 782(1997)69
Anderson, J.M., see Saari-Nordhaus, R. 782(1997)75
Arvidsson, T., see Sänger-van de Griend, C.E. 782(1997)271
Bartolini, P., see Dalmora, S. 782(1997)199
Başer, K.H.C., see Bozan, B. 782(1997)133
Beijnen, J.H., see Sparidans, R.W. 782(1997)211
Bielejewska, A., see Nowakowski, R. 782(1997)1
Blackwell, J.A., see Cantrell, G.O. 782(1997)237
Bozan, B., Başer, K.H.C. and Kara, S.
Quantitative determination of naphthaquinones of *Arnebia densiflora* (Nordm.) Ledeb. by an improved high-performance liquid chromatographic method 782(1997)133
Buzzi, M., see Stefani, R. 782(1997)123
Cantrell, G.O. and Blackwell, J.A.
Comparison of 1,1,1,2-tetrafluoroethane and carbon dioxide-based mobile phases for packed column supercritical fluid chromatography 782(1997)237
Capangpangan, M.B. and Suffet, I.H.(Mel)
Validation studies for a new supercritical fluid extraction method for the isolation of hydrophobic organic compounds from filtered suspended solids 782(1997)247
Carneiro, M.C., see Galceran, M.T. 782(1997)289
Cartoni, G.P., Coccioli, F. and Spagnoli, M.
Analysis of ethereal extracts of wines and other alcoholic beverages by high-performance liquid chromatography with microbore columns 782(1997)219
Castillo, G. and Martinez, S.
Reversed-phase C_{18} high-performance liquid chromatography of gibberellins GA_3 and GA_1 782(1997)137
Cavender, D.M., see Wolf, C. 782(1997)175
Coccioli, F., see Cartoni, G.P. 782(1997)219
Dalmora, S., de Oliveira, J.E., Affonso, R., Gimbo, E., Ribela, M.T.C.P. and Bartolini, P.
Analysis of recombinant human growth hormone directly in osmotic shock fluids 782(1997)199
De Keukeleire, D., see Nascimento, R.F. 782(1997)13
De Leenheer, A.P., see Huo, J.-Z. 782(1997)63
Den Hartigh, J., see Sparidans, R.W. 782(1997)211
Denton, G., see Murray, A. 782(1997)49
De Oliveira, J.E., see Dalmora, S. 782(1997)199
Derrico, E.M., see Wolf, C. 782(1997)175
Diez, M., see Galceran, M.T. 782(1997)289
Duszczyk, K., see Nowakowski, R. 782(1997)1
Elfakir, C. and Lafosse, M.
Porous graphitized carbon and octadecyl-silica columns in the separation of some alkylglycoside detergents 782(1997)191
Fanali, S., see Roos, N. 782(1997)257
Felici, M., see Gelsomino, A. 782(1997)105
Franco, D.W., see Nascimento, R.F. 782(1997)13
Galceran, M.T., Carneiro, M.C., Diez, M. and Puignou, L.
Separation of quaternary ammonium herbicides by capillary electrophoresis with indirect UV detection 782(1997)289
Ganzler, K., see Roos, N. 782(1997)257
Gelsomino, A., Petrovičová, B., Tiburtini, S., Magnani, E. and Felici, M.
Multiresidue analysis of pesticides in fruits and vegetables by gel permeation chromatography followed by gas chromatography with electron-capture and mass spectrometric detection 782(1997)105
Gimbo, E., see Dalmora, S. 782(1997)199
Grajek, H., Witkiewicz, Z. and Jankowska, H.
Application of Kováts retention indices for investigation of adsorption properties of activated carbons 782(1997)87
Grazzi, R., see Stefani, R. 782(1997)123
Gröningsson, K., see Sänger-van de Griend, C.E. 782(1997)271
Haddad, P.R., see Ohta, K. 782(1997)33
Haginaka, J. and Kanasugi, N.
Separation of basic drug enantiomers by capillary zone electrophoresis using ovoglycoprotein as a chiral selector 782(1997)281
Hayashi, Y., see Nishikawa, T. 782(1997)296
Helliger, W., see Lindner, H. 782(1997)55
Holm, T.
Mechanism of the flame ionization detector. II. Isotope effects and heteroatom effects 782(1997)81
Huo, J.-Z., Nelis, H.J., Lavens, P., Sorgeloos, P. and De Leenheer, A.P.
Determination of E vitamers in microalgae using high-performance liquid chromatography with fluorescence detection 782(1997)63
Jankowska, H., see Grajek, H. 782(1997)87
Kanasugi, N., see Haginaka, J. 782(1997)281
Kara, S., see Bozan, B. 782(1997)133
Kim, J.-H. and Lee, J.-H.
Simultaneous determination of six cations in mineral water by single-column ion chromatography 782(1997)140
Klingenberg, A., see Seubert, A. 782(1997)149
Konno, Y., see Kubota, N. 782(1997)159
Kubo, H., see Nishikawa, T. 782(1997)296
Kubota, N., Konno, Y., Saito, K., Sugita, K., Watanabe, K. and Sugo, T.
Module performance of anion-exchange porous hollow-fiber membranes for high-speed protein recovery 782(1997)159
Lafosse, M., see Elfakir, C. 782(1997)191
Lavens, P., see Huo, J.-Z. 782(1997)63
Lee, J.-H., see Kim, J.-H. 782(1997)140
Lima Neto, B.S., see Nascimento, R.F. 782(1997)13
Lin, J.-T., Woodruff, C.L. and McKeon, T.A.
Non-aqueous reversed-phase high-performance liquid chromatography of synthetic triacylglycerols and diacylglycerols 782(1997)41
Lindner, H., Sarg, B. and Helliger, W.
Application of hydrophilic-interaction liquid chromatography to the separation of phosphorylated H1 histones 782(1997)55
Magnani, E., see Gelsomino, A. 782(1997)105
Marques, J.C., see Nascimento, R.F. 782(1997)13

Martinez, S., see Castillo, G. 782(1997)137
Maruška, A. and Pyell, U.
Capillary electrochromatography: normal-phase mode using silica gel and cellulose-based packing materials 782(1997)167
McKeon, T.A., see Lin, J.-T. 782(1997)41
Montgomery, R.M., see Saari-Nordhaus, R. 782(1997)69
Moritz, T., see Wingsle, G. 782(1997)95
Mounts, T.L., see Abidi, S.L. 782(1997)25
Murray, A., Sekowski, M., Spencer, D.I.R., Denton, G. and Price, M.R.
Purification of monoclonal antibodies by epitope and mimotope affinity chromatography 782(1997)49
Nair, L.M., see Saari-Nordhaus, R. 782(1997)69
Nascimento, R.F., Marques, J.C., Lima Neto, B.S., De Keukeleire, D. and Franco, D.W.
Qualitative and quantitative high-performance liquid chromatographic analysis of aldehydes in Brazilian sugar cane spirits and other distilled alcoholic beverages 782(1997)13
Nelis, H.J., see Huo, J.-Z. 782(1997)63
Nishikawa, T., Hayashi, Y., Suzuki, S., Kubo, H. and Ohtani, H.
Erratum to "On-column enantiomerization of 3-hydroxybenzodiazepines during chiral liquid chromatography with optical rotation detection". [J. Chromatogr. A, 767 (1997) 93–100] 782(1997)296
Nowakowski, R., Bielejewska, A., Duszczyk, K. and Sybilska, D.
Chiral discrimination by high-performance liquid chromatography with joint use of two cyclodextrin additives 782(1997)1
Ohta, K., Tanaka, K. and Haddad, P.R.
Ion-exclusion chromatography of benzenecarboxylic acids on an unmodified silica-gel column 782(1997)33
Ohtani, H., see Nishikawa, T. 782(1997)296
Petrovičová, B., see Gelsomino, A. 782(1997)105
Pirkle, W.H., see Wolf, C. 782(1997)175
Price, M.R., see Murray, A. 782(1997)49
Puignou, L., see Galceran, M.T. 782(1997)289
Purdy, W.C., see Spencer, B.J. 782(1997)227
Pyell, U., see Maruška, A. 782(1997)167
Ribela, M.T.C.P., see Dalmora, S. 782(1997)199
Roos, N., Ganzler, K., Szemán, J. and Fanali, S.
Systematic approach to cost- and time-effective method development with a starter kit for chiral separations by capillary electrophoresis 782(1997)257
Rozing, G.P., see Wolf, C. 782(1997)175
Saari-Nordhaus, R. and Anderson, Jr, J.M.
Electrochemically regenerated ion suppressors, new suppressors for ion chromatography 782(1997)75
Saari-Nordhaus, R., Nair, L.M., Montgomery, R.M. and Anderson, Jr, J.M.
Electro-elution ion chromatography: controlling sample retention by electrochemically generating or modifying the mobile phase 782(1997)69
Saito, K., see Kubota, N. 782(1997)159
Sänger-van de Griend, C.E., Gröningsson, K. and Arvidsson, T.
Enantiomeric separation of a tetrapeptide with cyclodextrin. Extension of the model for chiral capillary electrophoresis by complex formation of one enantiomer molecule with more than one chiral selector molecules 782(1997)271
Sarg, B., see Lindner, H. 782(1997)55
Sekowski, M., see Murray, A. 782(1997)49
Seubert, A. and Klingenberg, A.
Sulfoacylated macroporous polystyrene–divinylbenzene: a new type of cation exchanger for the analysis of multivalent metal cations 782(1997)149
Sorgeloos, P., see Huo, J.-Z. 782(1997)63
Spagnoli, M., see Cartoni, G.P. 782(1997)219
Sparidans, R.W., den Hartigh, J., Beijnen, J.H. and Vermeij, P.
Derivatization of pamidronate and other amino(bis)phosphonates with different isothiocyanates prior to ion-pair liquid chromatography 782(1997)211
Spence, P.L., see Wolf, C. 782(1997)175
Spencer, B.J. and Purdy, W.C.
Comparison of the separation of fat-soluble vitamins using β-cyclodextrins in high-performance liquid chromatography and micellar electrokinetic chromatography 782(1997)227
Spencer, D.I.R., see Murray, A. 782(1997)49
Stefani, R., Buzzi, M. and Grazzi, R.
Supercritical fluid extraction of pesticide residues in fortified apple matrices 782(1997)123
Suffet, I.H.(Mel), see Capangpangan, M.B. 782(1997)247
Sugita, K., see Kubota, N. 782(1997)159
Sugo, T., see Kubota, N. 782(1997)159
Suzuki, S., see Nishikawa, T. 782(1997)296
Sybilska, D., see Nowakowski, R. 782(1997)1
Szemán, J., see Roos, N. 782(1997)257
Tanaka, K., see Ohta, K. 782(1997)33
Tiburtini, S., see Gelsomino, A. 782(1997)105
Vermeij, P., see Sparidans, R.W. 782(1997)211
Wan, Q.-H.
Capillary electrochromatography: effect of electrolyte concentration on electroosmotic flow and column efficiency 782(1997)181
Watanabe, K., see Kubota, N. 782(1997)159
Wingsle, G. and Moritz, T.
Analysis of ascorbate and dehydroascorbate in plant extracts by high-resolution selected ion monitoring gas chromatography–mass spectrometry 782(1997)95
Witkiewicz, Z., see Grajek, H. 782(1997)87
Wolf, C., Spence, P.L., Pirkle, W.H., Derrico, E.M., Cavender, D.M. and Rozing, G.P.
Enantioseparations by electrochromatography with packed capillaries 782(1997)175
Woodruff, C.L., see Lin, J.-T. 782(1997)41